

Angular Dependences of Third Harmonic Generation from Microdroplets

J. Kasparian^(a), B. Krämer^(b), J. P. Dewitz^{(c)2}, S. Vajda^(b), P. Rairoux^{(b)1}, B. Vezin^(a), V. Boutou^(a), T. Leisner^(b),
W. Hübner^(c), J. P. Wolf^(a), L. Wöste^(b), K. H. Bennemann^(c)

(a) LASIM (UMR 5579), Université Claude Bernard Lyon 1, 43 bd du 11 novembre, 69622 Villeurbanne Cedex France

(b) Institut für Experimentalphysik, Freie Universität Berlin, Arnimallee 14, 14195 Berlin, Germany

(c) Institut für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, 14195 Berlin, Germany.

(February 1, 2008)

We present experimental and theoretical results for the angular dependence of third harmonic generation (THG) of water droplets in the micrometer range (size parameter $62 < ka < 248$). The THG signal in *p*- and *s*-polarization obtained with ultrashort laser pulses is compared with a recently developed nonlinear extension of classical Mie theory including multipoles of order $l \leq 250$. Both theory and experiment yield over a wide range of size parameters remarkably stable intensity maxima close to the forward and backward direction at “magic angles”. In contrast to linear Mie scattering, both are of comparable intensity.

42.65.Ky;78.40.Dw;42.68.Ge;92.60.Jq

Studies of optical microcavities have received increasing interest both from fundamental [1] as well as applied research [2,3]. The quality factor of microcavities formed by liquid droplets can become as large as 10^6 . This leads to very high internal electromagnetic fields under resonant excitation. Accordingly, nonlinear optical effects including stimulated Raman scattering (SRS), third order sum frequency generation (TOSF) and in particular third harmonic generation (THG) in microparticles are enhanced by several orders of magnitude compared to flat surfaces and bulk media [4,5].

These enhanced effects, in turn, make nonlinear optics a promising spectroscopic probe for the characterization of microparticles. With respect to applications in this area a lot of interest is focused on determining size, shape, and refractive index from the measured intensities. With the help of the recently developed lasers generating ultra-short pulses of high intensities, higher harmonic generation can thus act as an additional source of information.

Optical TOSF in particular THG in droplets was first reported by Chang and coworkers in many organic and inorganic liquids [6,7]. Theoretical models for TOSF and angle integrated SHG in microdroplets have been proposed by several authors [8]. The angular dependence of SHG and THG by particles with size parameter $ka \leq 5$ ($ka = 2\pi a/\lambda$, a being the radius of the particle) has recently been investigated by Dewitz *et al.* using a nonlinear extension of classical Mie theory [9].

Here we report the first observation of the angular dependence of THG generated by micrometer-sized water droplets ranging from 8 to 32 μm ($62 < ka < 248$), as well as new calculated angular distributions for droplets with radius up to about 10.5 μm ($ka \approx 81$). Theory and experiment yield in fair agreement the following three remarkable features:

- The THG scattering profile of droplets having a radius

- of $a \gtrsim 8 \mu\text{m}$ is dominated by only a few intensity maxima at magic angles close to the forward and backward directions.

- The intensity and the angular distribution of the THG signal depends only weakly on the particle size and the central wavelength of the laser.

- Theory and experiment find comparable intensity close to the forward and backward direction.

Each of these findings is in strong contrast to linear Mie scattering and is of considerable relevance for the application of nonlinear optical spectroscopy to micro-particle sizing and detection.

Our theoretical model for higher harmonic generation by microdroplets is similar to the formalism of linear Mie theory [10]. Therein the electrodynamic response of spherical particles is described by matching the incident, internal and scattered field with the help of the boundary conditions of their parallel components. The expansion coefficients of the scattered field can then be expressed in terms of the coefficients of the incident field. On this basis Östling *et al.* [5] developed a model for angle-integrated higher harmonic generation by microparticles. This theory has been extended to angular dependences by three of the present authors [9]. The main idea of the approach is that in a first step the incident field generates a charge $\sigma = \sum_{n=1}^{\infty} \sigma^{(n)}$, where $\sigma^{(n)}$ oscillates with frequency $n\omega$. This picture is motivated by the anharmonic oscillator model [9]. In a second step the nonlinear components $\sigma^{(n)}$ radiate the n -th harmonic field. In the central approximation of the theory $\sigma^{(n)}$ is set equal to $[\sigma^{(1)}]^n$, where $\sigma^{(1)}$ is equal to the radial component of the polarization $\mathbf{P}^{(1)}$ known from linear Mie theory [5]. The n -th power of $\sigma^{(1)}$ can easily be obtained by expanding $\sigma^{(1)}$ in terms of spherical harmonics taking into account the appropriate selection rules for the angular momentum and keeping only terms varying with $e^{in\omega t}$. Using $\sigma^{(n)}$ in combination with the boundary conditions

$\mathbf{n} \cdot [\mathbf{D}_{\text{out}}^{(n)} - \mathbf{D}_{\text{in}}^{(n)}] = 4\pi\sigma^{(n)}$ for the radial components and $\mathbf{n} \times [\mathbf{E}_{\text{out}}^{(n)} - \mathbf{E}_{\text{in}}^{(n)}] = 0$ for the parallel components and the appropriate expansion of the fields in vector spherical harmonics [11], one determines then the radiated intensities from the radial part of the Poynting vector $|\mathbf{n} \cdot (\mathbf{E} \times \mathbf{H})|$. Here the subscripts “in” and “out” denote the fields inside and outside the droplet [12].

In THG multipoles of order $l \leq 3 \cdot ka + \delta$ ($\delta \approx 20$ according to the required precision) must be kept in the multipole expansions in order to get a fully converged series expansion for the fields. Although conceptionally straightforward it is numerically very time-consuming to obtain higher-order multipole contributions. Consequently our previous calculations were restricted to size parameters $ka \leq 5$. For *small* spherical droplets in this size range the theoretical treatment showed for SHG and THG an increasing number of maxima which were distributed over the whole angular profile. They result from an increased number of multipoles. The angles of the intensity maxima exhibited strong variations upon small changes of the droplet size parameter ka . Moreover, the enhancement of the intensities scattered in the forward direction -well known from linear Mie scattering [13,14]-, was amplified in SHG and THG dominating the angular profile. In the present paper we focus on larger particles ($ka > 62$) obtaining quite distinct results.

The experimental setup is given in Fig. 1. Monodisperse water droplets were generated at a repetition rate of 1 kHz by a piezo-driven nozzle. Depending on the operation condition of the nozzle, droplets in the size range between $8 \mu\text{m}$ and $32 \mu\text{m}$ could be produced. The droplet generation was synchronized to a Ti:Sa femtosecond laser system, which produces ultrashort light pulses (80 fs) with pulse energies up to $400 \mu\text{J}$ at a central wavelength of 820 nm. The scattered light was collected with a quartz fiber mounted onto a stepper motor driven goniometer, then spectrally dispersed into a UV spectrometer and detected by a photomultiplier tube. Polarizing beamsplitters could be inserted in front of the fiber to allow polarisation sensitive measurements. The setup allowed to analyze the radiation emitted from the droplet in an angular range between $18^\circ < \theta < 160^\circ$. The angular resolution was 1° . The shot to shot variation of the particle size was controlled independently by the analysis of the Mie scattering patterns obtained with a He:Ne laser and by direct observation of the droplets under a light microscope; it was found to be below 2 %. The stability of the droplet generation and the temporal and spatial overlap of the Ti:Sa laser beam and droplet stream was continuously monitored by observing the scattered laser light with a microscope which was equipped with a CCD camera readout. At higher fluences of the Ti:Sa laser, the droplets disintegrated into a cloud of small droplets with a diameter of about $1 \mu\text{m}$.

Since the optical response of the droplets to the ultra-

short laser pulses is much faster than their fragmentation, the light scattering signal was nevertheless detected. It was checked experimentally that background signal from remaining small aerosol droplets could be neglected compared to the initial droplet signal. In order to determine the dominant channel of the nonlinear optical response we performed spectroscopic measurements. We find as the most significant feature a large third harmonic peak at 273 nm. It showed a cubic dependence of the incident laser fluence, as expected for a third order process. The SHG peak at 410 nm is rather weak. This is an indication for the good spherical shape of the droplets for which SHG is forbidden by symmetry. The THG light was *s*- or *p*-polarized as the incident light, the amount of the cross polarization was only 2%. Moreover, no stimulated Raman scattering was observed. This is comprehensible with respect to the short laser pulses used and demonstrates that THG is indeed the dominant nonlinear response channel.

In Fig. 2 the experimental results for the angular dependence of THG of water droplets having a radius of $a = 10.5 \mu\text{m}$ ($ka \simeq 81$) are shown for angles between 18° and 160° in the $\varphi = 0^\circ$ plane. Two pronounced intensity maxima appear around 26° and 155° , whereas almost no intensity is detected at angles between 50° and 130° . The result is in clear contrast to the angular dependence expected for linear Mie scattering from droplets with $ka \approx 81$. Instead of a very complex structure with several maxima of varying angle and magnitude only one broad maximum appears close to the forward and backward directions.

Moreover we find that the angular dependence is remarkably stable upon variation of the droplet size: the scattering profile is reproduced for particles from $a = 8 \mu\text{m}$ to $32 \mu\text{m}$. Only a small shift of the “magic angle” of maximum intensity occurs from 26° to 30° . Thus the angular dependence does not only show a strong reduction of complexity but also a *stabilization of the angles* of intensity maxima over a wide range of droplet sizes, although the number of contributing multipoles varies strongly with ka .

Most importantly we find that the magnitude of the intensity maxima close to the forward and backward direction is of the same order. This is in sharp contrast to linear Mie scattering where forward scattering is strongly enhanced.

In Fig. 3 we show the theoretical results for the angular dependence of the THG intensities of a water droplet with radius $a = 10.5 \mu\text{m}$ ($ka \approx 81$) in the full θ -range from 0° to 180° , in the inset the corresponding polar plot is displayed. Dominating maxima appear close to the forward and backward direction at angles $\theta \approx 10^\circ$ (where no experimental data could be recorded) and $\theta \approx 158^\circ$. Those maxima show unexpected stability upon droplet size variation. The stability of intensity maxima in THG strongly contrasts with linear Mie scattering, where the-

ory yields a strong change of the scattering profile upon variation of ka in a small range around $ka \approx 81$ [16]. Furthermore the magnitudes of the intensity maxima close to the forward and backward directions are of same order in THG as it is observed in experiments (see Fig. 2). This is again in clear contrast to linear Mie scattering for which the calculated ratio of forward to backward scattering for water droplets with $ka \approx 81$ is around 6700 [16].

The stabilization of the angular scattering profile in THG as a function of ka results from dominating resonances occurring within the laser bandwidth. These sharp resonance peaks are enhanced by six orders of magnitude as compared to the background radiation. The dramatic enhancement results from the nonlinearity in THG, which amplifies the field effects. Remarkably, our calculations yield that every resonance displays the same angular pattern. Experimentally these resonances are always excited due to the finite laser bandwidth of ± 5 nm resulting from the femtosecond pulses and the particle size fluctuation of $\pm 2\%$. In our case these effects have been fully taken into account in the calculations. In the experimental ka -range ($ka = 81 \pm 2$) four resonances are excited.

In backward direction the position of the two peaks with the largest angles (156° , 160°) are found by both theory and experiment. For decreasing angle the agreement fades since the theory shows a more complicated structure and thus a stronger oscillation of the intensities. The damping of the experimental structure results from the finite angular resolution which is of the order of 1° . For intensities less than 130° both theory and experiment show no structure.

In the forward direction theory and experiment agree in a less satisfactory way. Calculations predict a strong maximum at an angle of 10° . This, however, is not observable with our existing equipment. Experimentally a maximum is observed at 26° . Work is in progress to investigate the angular range also below 18° .

In conclusion, the angular dependence of THG of water droplets with sizes from $8 \mu\text{m}$ up to $32 \mu\text{m}$ are studied experimentally and for sizes around 9 and $10 \mu\text{m}$ theoretically. In good agreement both investigations show a reduced complexity of the scattering profile and a stabilization of the angles of maximum intensities close to the forward and backward direction. Moreover the magnitude of the intensities in the forward and backward direction are of the same order of magnitude. These results are in remarkable contrast to the characteristics of linear Mie scattering in this size range.

The angles of the maxima near the backward direction show good quantitative agreement between theory and experiment. It was shown that the stabilization is an effect of the excitation of sharp resonance peaks with stable angular dependence independent of the size of the droplet.

This stabilization can be considered in analogy to rain-

bow scattering which occurs in linear optics for large droplet sizes and reflects the onset of a transition from Mie scattering to geometrical optics. The latter is valid for a plane surface resulting from the limit of droplets with infinite radius. Due to the incorporation of higher multipoles in THG this limit is reached for smaller droplets than in linear optics.

Thus THG opens up new possibilities to characterize particles in the Mie-size range between Rayleigh-scattering and geometrical optics. Especially the strongly enhanced backward scattering yields new perspectives for the remote sensing of the atmosphere.

J. P. Dewitz and W. Hübner gratefully acknowledge financial support through Deutsche Forschungsgemeinschaft Sfb 290. The authors of LASIM also acknowledge the DRET for actively supporting this project.

-
- ¹ Alfred Wegener Institut für Polar- und Meeresforschung, Telegraphenberg A 43, 14401 Potsdam, Germany
² e-mail: dewitz@physik.fu-berlin.de
- [1] J. C. Knight, N. Dubreuil, V. Sandoghdar, J. Harc, V. Lefèvre-Seguin, J. M. Raimond, S. Haroche, *Opt. Lett.* **20**, 1515 (1995)
 - [2] S. Arnold, N. Neumann, A. B. Pluchino, *Opt. Lett.* **9**, 4 (1984)
 - [3] G. S. Grader, S. Arnold, R. C. Flagan, J. H. Steinfeld, *J. Chem. Phys.* **86**, 5897 (1987)
 - [4] S. X. Qian, J. B. Snow, R. K. Chang, in: *Laser Spectroscopy VII*, T. W. Hänsch, Y. R. Shen (eds.), Springer Verlag Berlin, 1985), p.204
 - [5] D. Östling, P. Stampfli, K. H. Bennemann, *Z. Phys. D* **28**, 169 (1993)
 - [6] W. P. Acker, D. H. Leach, R. K. Chang, *Opt. Lett.* **14**, 402 (1989)
 - [7] D. H. Leach, R. K. Chang, W. P. Acker, S. C. Hill, *J. Opt. Soc. Am. B* **10**, 34 (1993)
 - [8] S. C. Hill, D. H. Leach, R. K. Chang, *J. Opt. Soc. Am.* **10**, 16 (1992); X. M. Hua, J. I. Gersten, *Phys. Rev. B* **33**, 3756 (1986); K. Hayata, M. Koshiba, *Phys. Rev. A* **46**, 6104 (1992)
 - [9] J. P. Dewitz, W. Hübner, K. H. Bennemann, *Z. Phys. D* **37**, 75 (1996)
 - [10] Mie G.: *Ann. Phys.* **25**, 377 (1908)
 - [11] Jackson J. D.: *Classical Electrodynamics* (2nd ed.), New York: Wiley 1975
 - [12] As a consequence of the approximation $\sigma^{(n)} \equiv [\sigma^{(1)}]^n$ the nonlinear susceptibility $\chi^{(n)}$ is taken to be constant and set equal to unity, thus the absolute value of the radiated intensities cannot be calculated. However, the theory still yields the full information about the angular dependence of the intensities and their variation with size, shape, and refractive index, since all interferences of the contributing multipoles are obtained.

- [13] Born M., Wolf E.: Principles of Optics, Oxford: Pergamon Press 1975
- [14] Bohren C. F., Huffman D. R.: Absorption and Scattering of Light by Small Particles, New York: Wiley 1983
- [15] Note however, that for ideally transparent objects interference effects similar to the shape resonances discussed in our paper cannot be neglected.
- [16] J. P. Dewitz, unpublished

FIG. 1. Experimental setup

FIG. 2. Measured spectra of THG intensity of a water droplet with $a = 10.5\mu\text{m}$ as a function of θ for parallel polarization. The forward and backward scattering is recorded separately as indicated by the dashed line. Thus the absolute intensity of both scans cannot be compared exactly.

FIG. 3. Theoretical spectra of the THG intensity of a water droplet with $a = 10.5\mu\text{m}$ in arbitrary units, represented linearly in the main part and as a polar plot in the inset. Parallel polarization is considered.